

Title	Fundamental Groups and Diophantine Geometry
Author(s)	Kim, Minhyong
Citation	代数幾何学シンポジウム記録 (2006), 2006: 116-118
Issue Date	2006
URL	http://hdl.handle.net/2433/214818
Right	
Type	Departmental Bulletin Paper
Textversion	publisher

Fundamental groups and Diophantine geometry

Minhyong Kim

January 17, 2007

Consider the strategy of studying a point x on a curve X (defined over \mathbb{Q}) by associating to it a macroscopic geometric object. There are at least three important examples that come to mind:

- (1) The Abel-Jacobi map that associates to x a line bundle $\mathcal{O}(x - b)$.
- (2) The Kodaira-Parshin construction that associates to x a curve C_x (running in a family parametrized by X).
- (3) The Frey-Hellegouarch construction that associates an elliptic curve to a point on a Fermat curve.

There is an obvious sense in which example (1) is much more canonical than the ad hoc methods of (2) and (3). However, (1) has never been generally applicable to finiteness theorems or the problem of determining points on hyperbolic curves.

The Abel-Jacobi map in the incarnation above is *algebraic*, as is important for Diophantine applications. Over the complex numbers, there is also an *analytic* description, whereby points of X parametrize extensions in the category of mixed Hodge structures:

$$\kappa_2^B : x \mapsto [0 \rightarrow H_1(X(\mathbb{C}), \mathbb{Z}) \rightarrow H_1(X(\mathbb{C}), \{x, b\}; \mathbb{Z}) \rightarrow \mathbb{Z} \rightarrow 0]$$

In fact, from this analytic perspective, Hain [5] constructed a lift of κ_2^B to a *higher Albanese map*:

$$\kappa^B : x \mapsto [P^B(X(\mathbb{C}); b, x)]$$

Here, $P^B(X(\mathbb{C}); b, x)$ is the set of unipotent paths from b to x :

$$P^B(X(\mathbb{C}); b, x) = \text{Isom}^\otimes(F_b, F_x)$$

where F_x is the fiber functor

$$F_x : \text{Un}(X(\mathbb{C}), \mathbb{Q}) \mapsto \text{Vect}_{\mathbb{Q}}$$

that associates to a local system \mathcal{L} its stalk at x :

$$\mathcal{L} \mapsto \mathcal{L}_x.$$

Thus, the $P^B(X(\mathbb{C}); b, x)$ for varying x runs through a classifying space for torsors under the pro-unipotent fundamental group

$$U^B(X(\mathbb{C}), b) := P^B(X(\mathbb{C}); b, b)$$

where the objects live in a suitable category of *non-abelian mixed Hodge structures* [6].

This construction in the realm of Archimedean analysis has a non-Archimedean analogue over \mathbb{Q}_p that gives rise to a map

$$\kappa^{dr/cr} : x \mapsto [P^{dr/cr}(X(\mathbb{Z}_p); b, x)]$$

going from \mathbb{Z}_p points of X to classifying spaces for torsors under the De Rham/crystalline fundamental group

$$U^{dr/cr}(X \otimes \mathbb{Q}_p, b).$$

Globally, there is also a *topological* analogue

$$\kappa^{et/glob} : x \mapsto [P^{et/glob}(\bar{X}; b, x)]$$

that takes global points $x \in X(R)$, where $R = \mathbb{Z}[1/S]$ is some ring of S -integers, to torsors under the unipotent étale fundamental group

$$U^{et/glob}(\bar{X}, b).$$

Recall that a motivic theory of such pro-unipotent fundamental groups was outlined by Deligne [1].

The eventual result of this discussion is a diagram:

$$\begin{array}{ccc} X(R) & \hookrightarrow & X(\mathbb{Z}_p) \\ \kappa^{et/glob} \downarrow & & \downarrow \kappa^{dr/cr} \\ H_f^1(\Gamma, U^{et}) & \xrightarrow{D \circ \text{loc}_p} & U^{dr}/F^0 \end{array}$$

The objects in the bottom row are pro-algebraic varieties over \mathbb{Q}_p that classify torsors in suitable categories. The bottom horizontal map comes from (easy) localization and (difficult) *non-abelian p -adic Hodge theory* ([3], [12]) which realizes the congruence between p -adic arithmetic topology and p -adic analysis. Eventually, we have an ambient pro-algebraic variety U^{dr}/F^0 and subsets $\text{Im}[X(\mathbb{Z}_p)]$ and $\text{Im}[H_f^1(\Gamma, U^{et})]$ that have the property of containing the image of the global points in their intersection:

$$\text{Im}[X(R)] \subset \text{Im}[X(\mathbb{Z}_p)] \cap \text{Im}[H_f^1(\Gamma, U^{et})]$$

The effort to use this intersection to gain control of the global points is described in the papers [2], [8], [9], [11], and [10]. The guiding principle that emerges is that the program outlined by Grothendieck in [4] should be viewed as a *nonabelian analogue* of the conjecture of Birch and Swinnerton-dyer [7].

References

- [1] Deligne, Pierre Le groupe fondamental de la droite projective moins trois points. Galois groups over \mathbb{Q} (Berkeley, CA, 1987), 79–297, Math. Sci. Res. Inst. Publ., 16, Springer, New York, 1989.
- [2] Faltings, Gerd Mathematics around Kim’s new proof of Siegel’s theorem. Preprint to be published in the proceedings of di Giorgi conference on number theory.
- [3] Fontaine, Jean-Marc Sur certains types de représentations p -adiques du groupe de Galois d’un corps local; construction d’un anneau de Barsotti-Tate. Ann. of Math. (2) 115 (1982), no. 3, 529–577.
- [4] Grothendieck, Alexandre Brief an G. Faltings, *Geometric Galois Actions, 1*, 49–58, London Math. Soc. Lecture Note Ser., 242, Cambridge University Press, Cambridge, 1997.
- [5] Hain, Richard M. Higher Albanese manifolds. Hodge theory (Sant Cugat, 1985), 84–91, Lecture Notes in Math., 1246, Springer, Berlin, 1987.
- [6] Hain, Richard M. The de Rham homotopy theory of complex algebraic varieties. I. *K-Theory* 1 (1987), no. 3, 271–324.
- [7] Kato, Kazuya p -adic Hodge theory and values of zeta functions of modular forms. Cohomologies p -adiques et applications arithmétiques. III. *Astrisque* No. 295 (2004), ix, 117–290.
- [8] Kim, Minhyong The motivic fundamental group of $\mathbb{P}^1 \setminus \{0, 1, \infty\}$ and the theorem of Siegel. *Invent. Math.* 161 (2005), no. 3, 629–656.

- [9] Kim, Minhyong The unipotent Albanese map and Selmer varieties for curves. Mathematics preprint archive, math.NT/0510441
- [10] Kim, Minhyong Remark on fundamental groups and effective Diophantine methods for hyperbolic curves. Preprint (2006).
- [11] Kim, Minhyong; Tamagawa, Akio The l -component of the unipotent Albanese map. Mathematics preprint archive, math.NT/0611384.
- [12] Vologodsky, Vadim Hodge structure on the fundamental group and its application to p -adic integration. Mosc. Math. J. 3 (2003), no. 1, 205–247, 260.